

Are Diffuse High Energy Neutrinos and γ -rays from Starburst Galaxies Observable?

F.W. Stecker

NASA Goddard Space Flight Center, Greenbelt, MD, USA

Abstract

Loeb and Waxman have argued that high energy neutrinos from the decay of pions produced in interactions of cosmic rays with interstellar gas in starburst galaxies would be produced with a large enough flux to be observable. Their model is reexamined here and we obtain an upper limit to the diffuse neutrino flux from starburst galaxies. The upper limit obtained here is a factor of ~ 5 lower than the flux which they predict. Our predicted neutrino flux would be below the atmospheric neutrino foreground flux at energies below ~ 300 TeV and therefore would be unobservable. Compared with predicted fluxes from other extragalactic high energy neutrino sources, starburst neutrinos with $\sim$ PeV energies would have a flux considerably below that predicted for AGN models.

We also estimate an upper limit for the diffuse GeV γ -ray flux from starburst galaxies to be $\mathcal{O}(10^{-2})$ of the observed γ -ray background, much less than the component from unresolved blazars and more than an order of magnitude below the estimate of Thompson *et al.*

Key words: extragalactic neutrinos; starburst galaxies

1 Introduction

Interactions of cosmic-ray nuclei with interstellar gas nuclei in our galaxy produce secondary pions. The neutral pions then decay to produce most of the galactic γ -rays above 0.1 GeV which have been observed [1]; the decay of the charged pions produces galactic cosmic-ray neutrinos [2].

It was pointed out 30 years ago that the distribution of high energy γ -rays in our Galaxy is related to the distribution of molecular clouds and very young hot high-mass stars in OB associations which are short-lived and explode into supernovae [3,4]. This association between supernovae which are likely to produce cosmic rays and dense regions of molecular gas led to the scenario

where interactions between the gas and cosmic rays then produce the γ -rays *via* the decay of the π^0 mesons produced in these interactions. A natural implication then would be that galaxies which were undergoing a phase of extremely active star formation would be likely sources of high energy γ -rays. Such sources are known as “starburst galaxies”. Detailed model calculations have been made to predict γ -ray fluxes above 100 MeV for the starburst galaxies Arp 220 [5] and NGC 253 [6]. The predicted flux for Arp 220 is close to the sensitivity limit for the *GLAST* large area telescope whereas the predicted flux for NGC 253 is an order of magnitude higher than that sensitivity limit¹ and close to the upper limit obtained from the *EGRET* data [7].

Loeb and Waxman (LW) have suggested that such hadronic processes in starburst galaxies, involving cosmic ray interactions with interstellar gas followed by the decay of $\pi^\pm$'s, can produce, *in toto*, a large enough background of diffuse high energy neutrinos to be observable [8] with a very large neutrino detector such as *Icecube* [9]. LW then argue that radio observations of starburst galaxies imply a *lower limit* on the cumulative extragalactic neutrino flux from starburst galaxies which is within the sensitivity range of *Icecube*. This argument is examined here and the opposite conclusion is obtained. Indeed, we derive an *upper limit* to the cumulative high energy neutrino flux from starburst galaxies. The diffuse flux of GeV γ -rays from the same processes is found to be $\mathcal{O}(10^{-2})$ of the observed γ -ray background, much less than the component from unresolved blazars and more than an order of magnitude below the estimate of Thompson *et al.* [10].

2 Radio Emission and the Neutrino Flux from Starburst Galaxies

LW start with the observed synchrotron emission from starburst galaxies which is produced by relativistic electrons in these sources [11]. They then make the assumptions that (1) the presence of relativistic electrons in these sources implies the presence of relativistic protons, (2) the protons lose essentially all of their energy to pion production, and (3), a lower limit to the energy loss rate of the protons can then be obtained from the synchrotron radio flux by assuming that all of the electrons (and positrons) which are radiating are from pion decay.

Assumption (1) is a reasonable one which is supported by observations of cosmic rays in our own Milky Way galaxy. Assumption (3), *viz.*, the “lower limit” assumption depends on assumption (2). However, assumptions (2) and (3) can be questioned because (a) the synchrotron radiating electrons may be largely accelerated primaries rather than secondaries related to pion production and

¹ For information on *GLAST* see <http://www-glast.stanford.edu>

decay as is the case in our own Galaxy, and (b) the conditions in starburst galaxies are significantly different from those in our own galaxy. In particular, starburst galaxies exhibit strong “superwinds” [14]. Such winds have significant dynamical effects and may disrupt magnetic fields and drive protons out of these galaxies before they can lose all of their energy by interacting with interstellar gas nuclei to produce pions. In contrast, assumption (2) of LW assumes full trapping of relativistic nuclei in the disks of starburst galaxies to the point where they only lose energy in hadronic interactions with gas atoms. This is in stark contrast to the situation in our own galaxy. (See footnote 2.)

These two caveats call into serious question the argument that the radio data can provide a true lower limit on the cumulative diffuse flux of neutrinos from starburst galaxies. Let us, however, ignore them and consider that assumptions (1)-(3) are reasonable for obtaining an analytic *upper limit* for such a flux. Let us then accept the other estimates which lead to the ratio of injected power of protons to electrons at a fixed particle energy, $\eta_{p/e} \simeq 6$ and a neutrino luminosity which is then related to the local radio luminosity density by

$$E_\nu^2 \Phi_\nu(E_\nu = 1 \text{ GeV}) \simeq (ct_H/4\pi)\zeta[4f(dL_f/dV)]_{f=1.4\text{GHz}} \quad (1)$$

where t_H is the age of the universe and $\zeta = 3$ is an evolution factor which takes account of the fact that starburst galaxies were more numerous in the past [8]. LW take the local energy production rate per unit volume at a frequency $f = 1.4$ GHz to be $\simeq 10^{28.5} \text{ W Mpc}^{-3}$. Let us reexamine this value for $f(dL_f/dV)]_{f=1.4\text{GHz}}$.

The local 1.4 GHz energy production rate has been derived by LW by making use of an important connection between radio emission and far infrared (FIR) emission in galaxies given in the paper of Yun, Reddy and Condon (YRC) [11]. That paper uses the data on *IRAS* galaxies to derive the local infrared luminosity density at $60 \mu\text{m}$ to be $2.6 \times 10^7 L_\odot \text{ Mpc}^{-3}$. This *total* power density is then used by LW to obtain the 1.4 GHz power density *via* a strong empirical correlation between the FIR and 1.4 GHz luminosity densities.

The key difference between the result to be derived here and that obtained by LW is in choosing how to interpret the paper of YRC. YRC state that less than 10% of the local FIR luminosity density is contributed by luminous IR galaxies with $L_{\text{FIR}} > 10^{11} M_\odot$; this is the component which includes the starburst galaxies. (Figure 11 of YRC yields an estimate of $\sim 6\%$.)

If we take the local contribution from starburst galaxies at $60\mu\text{m}$ to be $2.6 \times 10^6 L_\odot$ (the 10% upper limit found by YRC), using the relation (4) of YRC as shown in their Fig. 5(a) one finds that the component of the local radio luminosity density related to the starburst galaxies is at most $\Phi_{1.4\text{GHz}} = 10^{18.4} \text{ W Hz}^{-1} \text{ Mpc}^{-3}$ which, when multiplied by 1.4 GHz, gives $\Phi_{1.4\text{GHz}} < 10^{27.5} \text{ W}$

Mpc^{-3} . This value is an order of magnitude lower than the flux obtained by LW [8].²

This is not the whole story because there is a higher relative fraction of the energy input from the higher relative number of starburst galaxies at higher redshifts. To estimate this effect, we assume that the FIR background is proportional to the integrated star formation activity rate. The fraction of the FIR background, $\kappa(\Delta z)$ contributed by galaxies in different redshift ranges, Δz , is obtained from Ref. [12]. Then we multiply $\kappa(\Delta z)$ by the fraction of the FIR background contributed by starburst galaxies in different redshift ranges, $\xi(\Delta z)$, to estimate the mean fraction of the total FIR background contributed by starburst galaxies. Estimates for κ and ξ are shown in Table 1.

Table 1: Relative contributions to the ν Starburst Galaxy Flux (see text).

Redshift Range (Δz)	$\kappa(\Delta z)$ [12]	$\xi(\Delta z)$	Reference for ξ
0 to 0.2	10%	$< 10\%$	[11]
0.2 to 1.2	68%	$\sim 13\%$	[12]
> 1.2	22%	$\sim 60\%$	[13]

Using the results from Table 1, we estimate that 23% of the observed FIR background integrated over redshift is from starburst galaxies.

3 Observability of High Energy Neutrinos from Starburst Galaxies

The upper limit on the radio flux from starburst galaxies obtained above can be used to obtain an upper limit on the neutrino flux from starburst galaxies by using equation (1) as derived by LW. One then finds that the neutrino background energy flux from starburst galaxies would be at most $\sim 2 \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$. Such a flux would be undetectable above the atmospheric

² One might also ask why not consider more “normal” galaxies with lower FIR luminosities and add them all in to estimate a higher neutrino flux? We note that in the case of our own galaxy, cosmic rays lose only a small fraction of their energy in pion producing interactions before escaping the disk, contrary to assumption (2). Also, cosmic rays with energies above 1 PeV (the relevant range for producing 100 TeV neutrinos) have a differential particle spectrum $\propto E^{-3.2}$, much steeper than the E^{-2} spectrum assumed by LW. This spectrum probably reflects both a steeper composite source spectrum and a shorter confinement time than those at lower energies.

background neutrino flux, even if equation (1) is assumed to be valid when extrapolated to 300 TeV and even granting all of the assumptions made by LW.

Table 2: Neutrino Energy Fluxes ($\text{GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$)

ν Source	$E^2\Phi(10\text{TeV})$	$E^2\Phi(100\text{TeV})$	$E^2\Phi(1\text{PeV})$	Reference
Atm: AMANDA-II	2×10^{-6}	7×10^{-8}	$< 3 \times 10^{-9}$	[15]
Atm (Vertical)	7×10^{-7}	$\sim 2 \times 10^{-9}$	—	[16]
AMANDA-II Diff.Lim.	9×10^{-8}	9×10^{-8}	9×10^{-8}	[17]
Starburst Galaxies	$< 2 \times 10^{-8}$	$< 2 \times 10^{-8}$	$< 2 \times 10^{-8}$	This paper
AGN Cores	5×10^{-10}	10^{-8}	10^{-7}	[18]
AGN	3×10^{-9}	3×10^{-8}	2×10^{-7}	[19]
GRB	5×10^{-10}	3×10^{-9}	3×10^{-9}	[20]
Icecube Sensitivity	—	4×10^{-9}	4×10^{-9}	[21]

Table 2 shows a comparison of the upper limit on the flux from starburst galaxies given here with the atmospheric neutrino flux and with approximate model predictions of neutrino fluxes γ -ray bursts (GRB) and active galactic nuclei (AGN) along with detector array sensitivities. It can be seen from this table that at 100 TeV none of the extragalactic sources proposed will dominate over the atmospheric foreground. The table shows that the present preliminary upper limit on the diffuse neutrino energy flux below 1 PeV from AMANDA-II is $\sim 8.8 \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ in the 10 TeV to 1 PeV energy range [17]. The full *Icecube* detector array is expected to push down to a sensitivity of $\sim 10^{-9} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ in the energy range $100 \text{ TeV} < E_\nu < 100 \text{ PeV}$ after several years of observation.³ Under the extreme assumption that the primary cosmic ray spectra in all starburst galaxies are as hard as E^{-2} up to energies $\mathcal{O}(10 \text{ PeV})$, PeV neutrinos from starburst galaxies may be detectable just above the projected sensitivity of *Icecube*.⁴ However, as can be seen from Table 2, above 1 PeV the AGN models predict fluxes which will be significantly larger than the the atmospheric foreground (expected to be $< 3 \times 10^{-9}$) $\text{GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ at 1 PeV), as well as the fluxes from γ -ray bursts (GRB) and starburst galaxies.

³ F. Halzen, private communication

⁴ Even if we make a second extreme assumption that 100% of the IR galaxies at redshifts greater than 1.2 are starburst galaxies, we would still predict a neutrino flux $< 3 \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$.

4 Observability of Diffuse γ -rays from Starburst Galaxies

In a follow-up paper to LW, Thompson *et al.* [10] have estimated the contribution of π^0 -decay γ -rays from starburst galaxies to the observed γ -ray background in the GeV energy range. Using an E^{-2} primary spectrum they get estimates of $E^2\Phi(E)$ fluxes for both γ -rays and neutrinos of 3×10^{-7} GeV cm $^{-2}$ s $^{-1}$ sr $^{-1}$. This is a factor of 3 larger than the neutrino flux of LW because they estimate that all of the electrons in these galaxies are from $\pi^\pm$ decay and they make the further hypothesis that a significant fraction the synchrotron radiating electrons which emit at 1.4 GHz lose energy by processes other than synchrotron radiation (*viz.*, bremsstrahlung and ionization in dense gas clouds.) Should the diffuse differential neutrino particle spectrum continue $\propto E^{-2}$ up to 10 TeV and above, the flux predicted in Ref. [10] would be more than a factor of 3 above the AMANDA-II limit, although no such claim is made in Ref. [10].

Our estimated γ -ray flux for the same E^{-2} primary spectrum is $\sim 2 \times 10^{-8}$ GeV cm $^{-2}$ s $^{-1}$ sr $^{-1}$ for γ -ray energies less than ~ 10 GeV. This is $\mathcal{O}(10^{-2})$ of the observed flux of $\sim 1.4 \times 10^{-6}$ GeV cm $^{-2}$ s $^{-1}$ sr $^{-1}$ determined by the *EGRET* group [22]. Above ~ 10 GeV the background spectrum will steepen owing to absorption from pair production interactions with the extragalactic ultraviolet background radiation [23]. It should be noted in this context that Stecker and Salamon have shown that the bulk of the observed background can be produced by unresolved blazars [23]. Thus, the contribution to the diffuse γ -ray background from starburst galaxies should be unobservable. In this context, we note that almost all of the observed extragalactic GeV γ -ray sources are blazars; no starburst galaxies have been observed.

Acknowledgments

The author would like to thank Eliot Quataert for his comments and for bringing Ref. [10] to his attention. This work was supported by NASA Grant ATP03-0000-0057.

References

- [1] Stecker, F. W. 1977, *Astrophys. J.* **212**, 60.
- [2] Stecker, F. W. 1979, *Astrophys. J.* **228**, 919.
- [3] Stecker, F. W. 1975, *Phys. Rev. Letters* **35**, 188.

- [4] Stecker, F. W. 1976, *Nature* **260**, 412.
- [5] Torres, D. F. 2004, *Astrophys. J.* **617**, 966.
- [6] Domingo-Santamaría, E. and Torres, D. F. 2005, *Astron. and Astrophys.* **444**, 403.
- [7] Blom, J. J. *et al.* 1999, *Astrophys. J.* **516**, 744.
- [8] Loeb, A. and Waxman, E. *JCAP* 05(2006)003.
- [9] Halzen, F. 2006, e-print astro-ph/ 0602132.
- [10] Thompson, T. A., Quataert, E. and Waxman, E. 2006, e-print astro-ph/ 0606665.
- [11] Yun, M. S., Reddy, N. A. and Condon, J. J. 2001, *Astrophys. J.* **554**, 803.
- [12] Lagache, G., Dole, H. and Puget, J.-L. 2005, in *The Fabulous Destiny of Galaxies: Bridging Past and Present*, in press, e-print astro-ph/ 0509556.
- [13] Erb, D. K., *et al.* 2006, *Astrophys. J.* **646**, 107.
- [14] Hoopes, C. G. *et al.* 2003, *Astrophys. J.* **596**, L175.
- [15] Bouchta, A. 2006, e-print astro-ph/ 0606235.
- [16] Gaisser, T. K. 2005, *Phys. Scripta* **T121**, 51.
- [17] Hill, G.C. (for the IceCube Collaboration) 2006, paper presented at the *Neutrino 2006 Intl. Conf.*, Santa Fe, NM.
- [18] Stecker, F. W. 2005, *Phys. Rev. D* **72** 107301.
- [19] Mannheim, K., Protheroe, R. J. and Rachen, J. P. 2001, *Phys. Rev. D* **63** 023003.
- [20] Waxman, E. and Bahcall, J. N. 1997, *Phys. Rev. Letters* **78**, 2292.
- [21] The Icecube Collaboration (M. Ribordy *et al.*), e-print astro-ph/ 0509322.
- [22] Sreekumar, P. *et al* 1998, *Astrophys. J.* **494**, 523.
- [23] Stecker, F. W. and Salamon, M. H. 1996, *Astrophys. J.* **464**, 600.